

GPO PRICE \$ \_\_\_\_\_

CFSTI PRICE(S) \$ \_\_\_\_\_

Hard copy (HC) \$1,00

Microfiche (MF) ,50

#853 July 85

THE GROUP OF PERMUTATIONS ON 3 LETTERS  
AND THE PROPER ROTATIONS OF A  
3-DIMENSIONAL ORTHOGONAL FRAME

by

Albert Gluckman

August 3, 1965

FACILITY FORM 802

**N66 27029**  
(ACCESSION NUMBER)

6  
(PAGES)

TMX-56895  
(NASA CR OR TMX OR AD NUMBER)

\_\_\_\_\_  
(THRU)

1  
(CODE)

19  
(CATEGORY)

Goddard Space Flight Center  
Greenbelt, Maryland

# THE GROUP OF PERMUTATIONS ON 3 LETTERS AND THE PROPER ROTATIONS OF A 3-DIMENSIONAL ORTHOGONAL FRAME

The ensuing mathematical discussion concerning the group of permutations on 3 letters and the proper rotations of a 3-dimensional orthogonal Frame is a study of relationships existing between the group of permutations on 3 letters,  $\mathfrak{S}_3$ , and the 6 different elements of the unimodular group  $SL(3)$  which are products of 3 unimodular elements of the  $3!$  products formed by permuting the order of the multiplication of these 3 proper rotation matrices; where  $M_1$ ,  $M_2$ , and  $M_3$  describe rotation of the Frame  $\mathfrak{F}_3$  about the x-, y-, and z-coordinate axis of  $\mathfrak{F}_3$ . Rotation shall not be constrained to the planes formed by the coordinate axes of  $\mathfrak{F}_3$ . This condition is equivalent to a choice of angles of rotation in such wise, that no one of the unimodular elements  $M_1$ ,  $M_2$ , and  $M_3$  is an identity. The angles  $\theta_1$ ,  $\theta_2$ , and  $\theta_3$  of rotation about the x-, y-, and z-coordinate axis respectively are elements of the field  $\mathbb{R}^\#$  of real numbers.  $\mathfrak{F}_3$  itself is embedded in a 3-dimensional continuum.

Consider the set of proper rotation matrices  $\{M_1, M_2, M_3\}$  describing rotations of an orthogonal Frame  $\mathfrak{F}_3$  embedded in a 3-dimensional continuum about the x-, y-, and z-coordinate axes of  $\mathfrak{F}_3$ , respectively. The explicit representation of these matrices are

$$M_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \theta_1 & -\sin \theta_1 \\ 0 & \sin \theta_1 & \cos \theta_1 \end{bmatrix}, \quad M_2 = \begin{bmatrix} \cos \theta_2 & 0 & \sin \theta_2 \\ 0 & 1 & 0 \\ -\sin \theta_2 & 0 & \cos \theta_2 \end{bmatrix}, \quad M_3 = \begin{bmatrix} \cos \theta_3 & -\sin \theta_3 & 0 \\ \sin \theta_3 & \cos \theta_3 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

These unimodular matrices may of course, be multiplied together in 6 different order arrangements to form 6 different products ~ to the set of permutations on 3 letters.

I shall let  $M_1 \sim 1$ ,  $M_2 \sim 2$ , and  $M_3 \sim 3$ , and list the ordering of the 6 multiplications of these matrices as

1.  $M_1 M_2 M_3 \sim 123 \sim \sigma_1$ ,    4.  $M_2 M_3 M_1 \sim 231 \sim \sigma_4$
2.  $M_1 M_3 M_2 \sim 132 \sim \sigma_2$ ,    5.  $M_3 M_1 M_2 \sim 312 \sim \sigma_5$
3.  $M_2 M_1 M_3 \sim 213 \sim \sigma_3$ ,    6.  $M_3 M_2 M_1 \sim 321 \sim \sigma_6$

After computation of the corresponding traces of the matrix group elements, I find that

$$\cos \theta_2 \cos \theta_3 + \cos \theta_1 \cos \theta_2 + \cos \theta_1 \cos \theta_3 - \sin \theta_1 \sin \theta_2 \sin \theta_3 = \psi_- =$$

$$\text{tr}(M_1 M_2 M_3) = \text{tr}(M_2 M_3 M_1) = \text{tr}(M_3 M_1 M_2), \text{ and}$$

$$\cos \theta_2 \cos \theta_3 + \cos \theta_1 \cos \theta_2 + \cos \theta_1 \cos \theta_3 + \sin \theta_1 \sin \theta_2 \sin \theta_3 = \psi_+ =$$

$$\text{tr}(M_1 M_3 M_2) = \text{tr}(M_2 M_1 M_3) = \text{tr}(M_3 M_2 M_1)$$

The permutational arrangement of unimodular group products  $\sim$  to the trace  $\psi_- \sim$  to a cyclic permutation of letters from left to right; and the cyclical (or permutational) arrangement of unimodular group products  $\sim$  to  $\psi_+ \sim$  to a cyclic permutation of letters from right to left. The array of cyclic permutations  $\sim \psi_+$  may be achieved to by an initial inversion of 2 letters from the array of cyclic permutations  $\sim \psi_-$ .

Again, suppose I let  $M_1 \sim 1$ ,  $M_2 \sim 2$ , and  $M_3 \sim 3$ , and consider the permutation group of order  $3!$  with respect to multiplication,  $\{\sigma_1, \sigma_2, \sigma_3, \sigma_4, \sigma_5, \sigma_6; \cdot\}$ ; letting the group elements be expressed as

$$\sigma_1 = 1, \sigma_2 = (23), \sigma_3 = (12), \sigma_4 = (123), \sigma_5 = (132), \sigma_6 = (13).$$

Then I can arrange the following correspondences,

$$\{\sigma_1, \sigma_4, \sigma_5\} \sim \text{tr}(M_1 M_2 M_3) = \text{tr}(M_2 M_3 M_1) = \text{tr}(M_3 M_1 M_2)$$

$$\{\sigma_2, \sigma_3, \sigma_6\} \sim \text{tr}(M_1 M_3 M_2) = \text{tr}(M_2 M_1 M_3) = \text{tr}(M_3 M_2 M_1)$$

where  $\{1, (123), (132)\} \sim \psi_-$  and  $\{(23), (12), (13)\} \sim \psi_+$  are the sets consisting of permutation elements corresponding to their respective traces. The inverses of the permutation elements may be expressed as

$$\sigma_1^{-1} = 1, \sigma_2^{-1} = (32), \sigma_3^{-1} = (21), \sigma_4^{-1} = (213), \sigma_5^{-1} = (312), \sigma_6^{-1} = (31).$$

I can now easily determine the permutation elements in terms of their inverses as a necessary step prior to formation of a classifying list of mutually conjugate group elements. Observe that  $1 \equiv 1 \Rightarrow \sigma_1^{-1} = \sigma_1$ ,  $(32) \equiv (23) \Rightarrow \sigma_2^{-1} = \sigma_2$ ,  $(21) \equiv (12) \Rightarrow \sigma_3^{-1} = \sigma_3$ ,  $(213) \equiv (132) \Rightarrow \sigma_4^{-1} = \sigma_5$ ,  $(312) \equiv (123) \Rightarrow \sigma_5^{-1} = \sigma_4$ , and  $(31) \equiv (13) \Rightarrow \sigma_6^{-1} = \sigma_6$ .

*A listing of the calculations of the mutually conjugate group elements of the symmetric group on 3 letters.*

$$\sigma_1^{-1}\sigma_1\sigma_1 = \sigma_1$$

$$\sigma_1^{-1}\sigma_4\sigma_1 = \sigma_4$$

$$\sigma_1^{-1}\sigma_2\sigma_1 = \sigma_2$$

$$\sigma_1^{-1}\sigma_5\sigma_1 = \sigma_5$$

$$\sigma_1^{-1}\sigma_3\sigma_1 = \sigma_3$$

$$\sigma_1^{-1}\sigma_6\sigma_1 = \sigma_6$$

$$\sigma_2^{-1}\sigma_1\sigma_2 = \sigma_1$$

$$\sigma_2^{-1}\sigma_4\sigma_2 = (32)(123)(23) = (132) = \sigma_5$$

$$\sigma_2^{-1}\sigma_2\sigma_2 = \sigma_2$$

$$\sigma_2^{-1}\sigma_5\sigma_2 = (32)(132)(23) = (123) = \sigma_4$$

$$\sigma_2^{-1}\sigma_3\sigma_2 = (32)(12)(23) = (13) = \sigma_6$$

$$\sigma_2^{-1}\sigma_6\sigma_2 = (32)(13)(23) = (12) = \sigma_3$$

$$\sigma_3^{-1}\sigma_1\sigma_3 = \sigma_1$$

$$\sigma_3^{-1}\sigma_4\sigma_3 = (21)(123)(12) = (132) = \sigma_5$$

$$\sigma_3^{-1}\sigma_2\sigma_3 = (21)(23)(12) = (13) = \sigma_6$$

$$\sigma_3^{-1}\sigma_5\sigma_3 = (21)(132)(12) = (123) = \sigma_4$$

$$\sigma_3^{-1}\sigma_3\sigma_3 = \sigma_3$$

$$\sigma_3^{-1}\sigma_6\sigma_3 = (21)(13)(12) = (23) = \sigma_2$$

$$\sigma_4^{-1}\sigma_1\sigma_4 = \sigma_1$$

$$\sigma_4^{-1}\sigma_4\sigma_4 = \sigma_4$$

$$\sigma_4^{-1}\sigma_2\sigma_4 = (213)(23)(123) = (13) = \sigma_6$$

$$\sigma_4^{-1}\sigma_5\sigma_4 = (213)(132)(123) = (132) = \sigma_5$$

$$\sigma_4^{-1}\sigma_3\sigma_4 = (213)(12)(123) = (23) = \sigma_2$$

$$\sigma_4^{-1}\sigma_6\sigma_4 = (213)(13)(123) = (12) = \sigma_3$$

$$\sigma_5^{-1}\sigma_1\sigma_5 = \sigma_1$$

$$\sigma_5^{-1}\sigma_4\sigma_5 = (312)(123)(132) = (123) = \sigma_4$$

$$\sigma_5^{-1}\sigma_2\sigma_5 = (312)(23)(132) = (12) = \sigma_3$$

$$\sigma_5^{-1}\sigma_5\sigma_5 = \sigma_5$$

$$\sigma_5^{-1}\sigma_3\sigma_5 = (312)(12)(132) = (13) = \sigma_6$$

$$\sigma_5^{-1}\sigma_6\sigma_5 = (312)(13)(132) = (23) = \sigma_2$$

$$\sigma_6^{-1}\sigma_1\sigma_6 = \sigma_1$$

$$\sigma_6^{-1}\sigma_4\sigma_6 = (31)(123)(13) = (132) = \sigma_5$$

$$\sigma_6^{-1}\sigma_2\sigma_6 = (31)(23)(13) = (12) = \sigma_3$$

$$\sigma_6^{-1}\sigma_5\sigma_6 = (31)(132)(13) = (123) = \sigma_4$$

$$\sigma_6^{-1}\sigma_3\sigma_6 = (31)(12)(13) = (23) = \sigma_2$$

$$\sigma_6^{-1}\sigma_6\sigma_6 = \sigma_6$$

A listing which consists of mutually conjugate group elements of the symmetric group in 3 letters.

$$1 = \sigma_1 \sim C_1 = \{1\}$$

$$(23) = \sigma_2 \sim C_{(23)} = \{(23), (13), (12)\}$$

$$(12) = \sigma_3 \sim C_{(12)} = \{(12), (13), (23)\}$$

$$(123) = \sigma_4 \sim C_{(123)} = \{(123), (132)\}$$

$$(132) = \sigma_5 \sim C_{(132)} = \{(132), (123)\}$$

$$(13) = \sigma_6 \sim C_{(13)} = \{(13), (12), (23)\}$$

PROPOSITION. If  $M_1, M_2$ , and  $M_3$  are proper rotations of a 3-dimensional orthogonal Frame  $\mathcal{B}$ , where these unimodular elements of the group  $SL(3)$  are multiplied together in 6 different permutational arrangements, the elements of which the mutually conjugate classes  $C_1, C_{(123)}$ , and  $C_{(132)}$ , consist of are inner automorphisms of the elements of the alternating group  $\mathcal{A}_3$ ; and the automorphisms of  $C_1, C_{(123)}$ , and  $C_{(132)}$ , correspond

to  $\psi_-$  where the permutations  $\sigma_1 \sim M_1 M_2 M_3$ ,  $\sigma_4 \sim M_2 M_3 M_1$ , and  $\sigma_5 \sim M_3 M_1 M_2$ . The group  $\{C_1, C_{(123)}, C_{(132)}, \dots\} \cong \mathcal{G}_3$ .

DEMONSTRATION. The permutation symbols could be considered as products of transpositions. It is then easy for me to determine which of the permutations are elements of the alternating group  $\mathcal{G}_3$ .

$$\sigma_1 = (1)(2)(3) = 1 \quad \sigma_4 = (123) = (12)(13)$$

$$\sigma_2 = (23) \quad \sigma_5 = (132) = (13)(12)$$

$$\sigma_3 = (12) \quad \sigma_6 = (13)$$

$\therefore \{\sigma_1, \sigma_4, \sigma_5; \dots\} \equiv \mathcal{G}_3 \sim \text{tr}(M_1 M_2 M_3) = \text{tr}(M_2 M_3 M_1) = \text{tr}(M_3 M_1 M_2)$ . In other words, the alternating group  $\{1, (123), (132); \dots\} \equiv \mathcal{G}_3 \sim \psi_-$ . With respect to the group  $\mathcal{G}_3$ , the classes of the mutually conjugate elements are determined by calculation and correspondence to be:

$$C_1 = \{1\}$$

$$C_{(123)} = \{(123), (132)\}$$

$$C_{(132)} = \{(132), (123)\}$$

Thus the inner automorphisms  $= \sigma_1, = \sigma_4$ , and  $= \sigma_5$  of the classes of mutually conjugate elements,  $C_1$ ,  $C_{(123)}$ , and  $C_{(132)}$  respectively, correspond to  $\psi_- = \text{tr}(M_1 M_2 M_3) = \text{tr}(M_2 M_3 M_1) = \text{tr}(M_3 M_1 M_2)$ , and the group  $\{C_1, C_{(123)}, C_{(132)}; \dots\} \cong \mathcal{G}_3$  as can be demonstrated by direct calculation.

Q.E.D.

#### BIBLIOGRAPHY

1. Edgar Dehn, *Algebraic equations*, New York, Dover Publications, Inc., 1960.
2. J. S. Lomont, *Applications of finite groups*, New York, Academic Press, 1959.